

Product Information

SEREXIS™ 200 Biopharma Grade Sample Kit

Defined FBS Replacement for Vaccine Production, with Lipid Mix 180
sterile-filtered

Cat. No. B-ZER-K1 consists of 1 x B-ZER-100ML & 1 x LIX-3ML

General Information

SEREXIS™ 200 is a defined serum alternative developed as a functional replacement to traditional fetal bovine serum (FBS) for a broad range of cell culture applications within Biopharma production workflows. It is designed to provide FBS-like performance under controlled and reproducible conditions.

SEREXIS™ 200 can be used as a direct replacement for FBS at standard serum concentrations and is intended for seamless integration into established workflows. It supports reliable cell growth, adherence, proliferation, viability, and morphology across multiple cell types, including A549, BHK-21, HeLa, MDBK, MDCK, Vero, primary immune cells, keratinocytes, and fibroblasts.

SEREXIS™ 200 combines extensive expertise in chemically defined media development with decades of experience in high-quality FBS manufacturing. Its formulation includes essential trace elements, defined proteins, selected animal-derived, highly purified protein fractions to support robust cell growth while maintaining controlled and reproducible culture conditions.

SEREXIS™ 200 is designed for highly regulated applications requiring advanced sourcing control, comprehensive documentation, and an elevated viral safety profile. All animal-derived components are sourced exclusively from fully traceable Australian origins and processed under controlled conditions, including gamma irradiation. Additional comprehensive viral testing further supports regulatory compliance and safety in vaccine manufacturing, biotech production, and other demanding biopharma workflows.

For cell types with increased lipid or fatty acid requirements, such as Hybridoma cultures, **SEREXIS™ 200** can be complemented with Lipid Mix 180 (Cat. No. LIX-10ML), a chemically defined lipid supplement supporting membrane synthesis, cellular metabolism, and proliferation.

SEREXIS™ 200 is the Biopharma Grade version of **SEREXIS™ 180**, developed for vaccine manufacturing, biotech production, and other demanding biopharma workflows.

Key Benefits

- High Performance without FBS for robust growth and stable morphology
- Designed and validated across multiple cell types including cell lines and primary cells
- Direct and seamless integration of **SEREXIS™ 200** in established workflows without adaptations
- Improved reproducibility due to the defined, homogeneous formulation
- Tailored for biopharmaceutical applications with highest safety, regulatory compliance, and comprehensive documentation
- Optional Lipid Mix 180 for demanding cell types

Product Information

SEREXIS™ 200 Biopharma Grade Sample Kit

Defined FBS Replacement for Vaccine Production, with Lipid Mix 180 sterile-filtered

Cat. No. B-ZER-K1 consists of 1 x B-ZER-100ML & 1 x LIX-3ML

Product Specifications

	SEREXIS™ 200 Biopharma Grade Defined FBS Replacement for Vaccine Production	Lipid Mix 180 Chemically Defined Concentrate
Cat. No.	B-ZER-500ML (500 ml), B-ZER-100ML (100 ml)	LIX-3ML (3 ml)
Appearance	Clear yellow liquid	Clear liquid
Storage and shelf life	<ul style="list-style-type: none"> Store at ≤ -15°C Protect from light! Avoid repeated freeze-thaw cycles. Preparation of aliquots recommended. 	<ul style="list-style-type: none"> Store at +2°C to +8°C Protect from light! Never freeze Lipid Mix 180 or supplemented end products
Working concentration	Use at 10% as a supplement in cell culture media or according to established FBS concentration	Use at 0.25% together with SEREXIS™ 200 as additional supplement in cell culture media
Thawing	Overnight at +2°C to +8°C	N.A.
Shipping conditions	Frozen (Dry ice)	Ambient

Instructions for Use

General Notes

- The standard working concentration of **SEREXIS™ 200** is 10%. Cell line specific adjustments of the final working concentration may increase the performance.
- FBS can typically be directly substituted with **SEREXIS™ 200** without previous stepwise adaptation.
- Performance may depend on culture conditions such as medium composition, seeding density, and process parameters.
- Avoid repeated freeze-thaw cycles, as this may negatively impact product performance. Aliquoting upon first thaw is recommended.
- Never freeze Lipid Mix 180.
- Do not freeze **SEREXIS™ 200**, when supplemented with Lipid Mix 180.
- SEREXIS™ 200** is compatible with established detachment workflows using trypsin. Using trypsin inhibitors is not required. For sensitive cell types, follow the detachment protocol in section "Cell Detachment & Trypsin Inactivation".
- Do not heat inactivate **SEREXIS™ 200**. Due to the defined formulation, heat inactivation is not required.
- Minimize exposure of Lipid Mix 180 to air, as the supplement contains unsaturated fatty acids that may undergo oxidation upon prolonged exposure.
- SEREXIS™ 200** can be used solely or together with Lipid Mix 180 (Cat. No. LIX-3ML). A detailed recommendation for testing strategy can be found in section "How to Use **SEREXIS™ 200** with Lipid Mix 180"

Product Information

SEREXIS™ 200 Biopharma Grade Sample Kit

Defined FBS Replacement for Vaccine Production, with Lipid Mix 180 sterile-filtered

Cat. No. B-ZER-K1 consists of 1 x B-ZER-100ML & 1 x LIX-3ML

Test Strategy for SEREXIS™ 200 & Lipid Mix 180

Depending on cell type and application, **SEREXIS™ 200** should be initially tested without Lipid Mix 180 (Approach 1). For cells that do not show sufficient proliferation when adhering to Approach 1 or for cells with increased demands for fatty acids, Approach 2 is recommended:

Test Strategy	Approach 1: SEREXIS™ 200	Approach 2: SEREXIS™ 200 & Lipid Mix 180
Recommended	Recommended as initial test strategy, as suitable for most cell types	Recommended, when cells cultured in Approach 1 show reduced growth Recommended for cells with increased lipid or fatty acid requirements, such as Hybridoma or insect cell culture
Supplementation		
Basal Medium	450 ml	450 ml
SEREXIS™ 200	50 ml (10% v/v)	50 ml (10% v/v)
Lipid Mix 180	None	1.25 ml (0.25% v/v)

- General Note: Most cell types do not require Lipid Mix 180 and, therefore, should be initially tested only with **SEREXIS™ 200** at 10% (v/v) working concentration without Lipid Mix 180
- Creating aliquots of **SEREXIS™ 200** is strongly recommended to avoid repeated freeze-thaw-cycles.
- Do not freeze Lipid Mix 180.
- For further technical support contact our experts at techservice@capricorn-scientific.com

Aliquotation of SEREXIS™ 200

1. Thaw **SEREXIS™ 200** in a water bath at 37°C while shaking or overnight at +2°C to +8°C
2. Homogenize **SEREXIS™ 200** by slowly shaking. Do not vortex.
3. Transfer **SEREXIS™ 200** to the laminar hood.
4. Aliquot **SEREXIS™ 200** to units of 50 ml following proper aseptic techniques.
5. Depending on the intended use, proceed as indicated below:
 - a. For long-term storage, freeze the **SEREXIS™ 200** aliquots again at ≤-15°C. Avoid repeated freeze-thaw cycles
 - b. For direct use, use 50 ml **SEREXIS™ 200** to supplement 450 ml of basal cell culture medium, such as DMEM High Glucose, with L-Glutamine (Cat. No. DMEM-HA). After supplementation, use the cell culture medium within 6-8 weeks and store in darkness.

Adaptation and Culturing

Cell Type	Most cell types in adherent and suspension cultures
Culture Flask Type	As established in standard protocols
Medium Volume	As established in standard protocols
Inoculation Cell Density	As established in standard protocols
Temperature	37°C
CO ₂ Concentration	5.0% or as established in standard protocols

Product Information

SEREXIS™ 200 Biopharma Grade Sample Kit

Defined FBS Replacement for Vaccine Production, with Lipid Mix 180 sterile-filtered

Cat. No. B-ZER-K1 consists of 1 x B-ZER-100ML & 1 x LIX-3ML

Note: Cultures can be adapted to cell culture media supplemented with **SEREXIS™ 200** or to cell culture media supplemented with **SEREXIS™ 200** and Lipid Mix 180

Direct Adaptation

Usually, for cells grown in serum supplemented medium or other serum-free medium little or no adaptation is needed and the cells may be directly transferred to **SEREXIS™ 200** supplemented media. Otherwise, follow the Instruction for Sequential Adaptation. It is advisable to keep a backup culture in the original media until cells have adapted.

Sequential Adaptation

- Sequential adaptation to media containing 10% **SEREXIS™ 200** should be tested in parallel to achieve optimal growth conditions depending on the cell type.
- Subculture cells into medium supplemented with a 1:4 ratio of **SEREXIS™ 200** and FBS, which typically corresponds 0.5 ml **SEREXIS™ 200** and 1.5 ml FBS per 20 ml of cell culture medium (see table below). During the adaptation procedure seed at twice the normal seeding density.
- Subculture cells when confluency reaches 70–90%. Subculture the cells in fresh pre-warmed medium supplemented with 1:4 ratio of **SEREXIS™ 200** and FBS. Once consistent cell growth with high viability has been achieved, passage cells into medium supplemented with 1:2 ratio of **SEREXIS™ 200** and FBS.
- Repeat step 2 of this procedure, stepwise increasing the ratio of **SEREXIS™ 200** to FBS 3:4 followed by 9:10 until the cells are subcultured into medium supplemented with 100% **SEREXIS™ 200**. Multiple passages at each step may be needed.
- Continue to monitor and passage cells until consistent growth with high viability is achieved. After several passages in medium supplemented with **SEREXIS™ 200**, the culture is adapted.

Step	Ratio	Basal Medium [ml]	FBS [ml]	SEREXIS™ 200 [ml]
1	1:4	9	0.75	0.25
2	1:2	9	0.5	0.5
3	3:4	9	0.25	0.75
4	9:10	9	0.1	0.9
5	1:1	9	0	1.0

Cell Detachment & Trypsin Inactivation

SEREXIS™ 200 is compatible with established detachment workflows using trypsin. Further reagents, such as trypsin inhibitors are not required. For sensitive cell types, use sensitive detachment reagents, such as Accutase (Cat. No. ACC-1B) or adhere to the detachment protocol below:

1. Check the cells for confluency under the microscope. Most cell types should be passaged, when they are ~ 70% – 80% confluent
2. Transfer the culture flask under a laminar hood and remove the medium
3. Rinse cells with PBS (Cat. No. PBS-1A) or another Ca²⁺ and Mg²⁺-free salt solution and completely remove the buffer.
4. Add enough trypsin (e.g., Cat. No. TRY-2B) or trypsin/EDTA (e.g., Cat. No. TRY-3B) to the culture flask to cover all cells

Product Information

SEREXIS™ 200 Biopharma Grade Sample Kit

Defined FBS Replacement for Vaccine Production, with Lipid Mix 180
sterile-filtered

Cat. No. B-ZER-K1 consists of 1 x B-ZER-100ML & 1 x LIX-3ML

5. Incubate at 37°C, or for more sensitive cultures, at room temperature or +2°C to +8°C until cells are detached. Check for cell detachment continuously as prolonged exhibition to trypsin can cause cellular damage.
6. Collect the cells in a sterile 15 ml reaction tube and immediately add 5 – 10 ml of fresh **SEREXIS™ 200**.
7. Centrifuge the cells for 5 min using standard centrifugation conditions appropriate for the respective cell type.
8. Remove the supernatant and gently resuspend the cell pellet in fresh cell culture medium supplemented with 10% **SEREXIS™ 200**, and optional with 0.25% Lipid Mix 180.
9. Seed the cells to a distinct density according to established protocols in a sterile flask fresh cell culture medium with **SEREXIS™ 200**, and optional with 0.25% Lipid Mix 180.

Cryopreservation

SEREXIS™ 200 is not suitable for cryopreservation. It is recommended to preserve cells in specialized serum-free cryopreservation media such as FreezeMe Two (Cat. No. FM2-F) or in DMSO-free cryopreservation media such as FreezeMe ZERO (Cat. No. FMZ-50ML), which support long-term storage and maintenance of genetic integrity.

Formulation

This formulation is our proprietary composition and has no counterparts either in its composition, or in its action.

Precautions and Disclaimer

This product is for research use and further manufacturing only.

Help Needed?

If you have any further questions regarding this product, please do not hesitate to contact our cell culture experts by email (techservice@capricorn-scientific.com) or phone (+49 6424 944640).